

FULLY DEGENERATE POLY-BERNOULLI NUMBERS AND POLYNOMIALS

DAE SAN KIM AND TAEKYUN KIM

ABSTRACT. In this paper, we introduce the new fully degenerate poly-Bernoulli numbers and polynomials and investigate some properties of these polynomials and numbers. From our properties, we derive some identities for the fully degenerate poly-Bernoulli numbers and polynomials.

1. INTRODUCTION

It is well known that the Bernoulli polynomials are defined by the generating function

$$(1.1) \quad \frac{t}{e^t - 1} e^{xt} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}, \quad (\text{see [1-21]}).$$

When $x = 0$, $B_n = B_n(0)$ are called the Bernoulli numbers. From (1.1), we note that

$$(1.2) \quad B_n(x) = \sum_{l=0}^n \binom{n}{l} B_l x^{n-l}, \quad (n \geq 0),$$

and

$$(1.3) \quad B_0 = 1, \quad B_n(1) - B_n = \delta_{1,n}, \quad (n \in \mathbb{N}), \quad (\text{see [1, 19]}),$$

where $\delta_{n,k}$ is the Kronecker's symbol.

In [4], L. Carlitz considered the degenerate Bernoulli polynomials which are given by the generating function

$$(1.4) \quad \frac{t}{(1 + \lambda t)^{\frac{1}{\lambda}} - 1} (1 + \lambda t)^{\frac{x}{\lambda}} = \sum_{n=0}^{\infty} \beta_{n,\lambda}(x) \frac{t^n}{n!}.$$

When $x = 0$, $\beta_{n,\lambda} = \beta_{n,\lambda}(0)$ are called the degenerate Bernoulli numbers. From (1.1) and (1.4), we note that

$$(1.5) \quad \frac{t}{e^t - 1} e^{xt} = \lim_{\lambda \rightarrow 0} \frac{t}{(1 + \lambda t)^{\frac{1}{\lambda}} - 1} (1 + \lambda t)^{\frac{x}{\lambda}} = \sum_{n=0}^{\infty} \lim_{\lambda \rightarrow 0} \beta_{n,\lambda}(x) \frac{t^n}{n!}.$$

Thus, by (1.5), we get

$$(1.6) \quad B_n(x) = \lim_{\lambda \rightarrow 0} \beta_{n,\lambda}(x), \quad (\text{see [4, 16]}).$$

2000 *Mathematics Subject Classification.* 11B75, 11B83, 05A19, 05A40.

Key words and phrases. Fully degenerate poly-Bernoulli polynomial, Fully degenerate poly-Bernoulli number, Umbral calculus.

By (1.4), we get

$$(1.7) \quad \sum_{n=0}^{\infty} (\beta_{m,\lambda}(1) - \beta_{m,\lambda}) \frac{t^n}{n!} = t,$$

and

$$(1.8) \quad \beta_{n,\lambda}(x) = \sum_{l=0}^n \binom{n}{l} \beta_{l,\lambda} \lambda^{n-l} \left(\frac{x}{\lambda}\right)_{n-l}.$$

From (1.7), we have

$$(1.9) \quad \beta_{n,\lambda}(1) - \beta_{n,\lambda} = \delta_{1,n}, \quad (n \geq 0), \quad \beta_{0,\lambda} = 1.$$

Now, we consider the degenerate Bernoulli polynomials which are different from the degenerate Bernoulli polynomials of L. Carlitz as follows:

$$(1.10) \quad \frac{\log(1 + \lambda t)^{\frac{1}{\lambda}}}{(1 + \lambda t)^{\frac{1}{\lambda}} - 1} (1 + \lambda t)^{\frac{x}{\lambda}} = \sum_{n=0}^{\infty} b_{n,\lambda}(x) \frac{t^n}{n!}.$$

When $x = 0$, $b_{n,\lambda} = b_{n,\lambda}(0)$ are called the degenerate Bernoulli numbers.

Note. The degenerate Bernoulli polynomials are also called Daehee polynomials with λ -parameter (see [14]).

From (1.10), we note that

$$(1.11) \quad \begin{aligned} \frac{t}{e^t - 1} e^{xt} &= \lim_{\lambda \rightarrow 0} \frac{\log(1 + \lambda t)^{\frac{1}{\lambda}}}{(1 + \lambda t)^{\frac{1}{\lambda}} - 1} (1 + \lambda t)^{\frac{x}{\lambda}} \\ &= \sum_{n=0}^{\infty} \lim_{\lambda \rightarrow 0} b_{n,\lambda}(x) \frac{t^n}{n!}. \end{aligned}$$

By (1.1) and (1.11), we see that

$$B_n(x) = \lim_{\lambda \rightarrow 0} b_{n,\lambda}(x), \quad (n \geq 0).$$

The classical polylogarithm function $\text{Li}_k(x)$ is defined by

$$(1.12) \quad \text{Li}_k(t) = \sum_{n=1}^{\infty} \frac{t^n}{n^k}, \quad (k \in \mathbb{Z}), \quad (\text{see [11, 12]}).$$

It is known that the poly-Bernoulli polynomials are defined by the generating function

$$(1.13) \quad \frac{\text{Li}_k(1 - e^{-t})}{1 - e^{-t}} e^{xt} = \sum_{n=0}^{\infty} B_n^{(k)}(x) \frac{t^n}{n!}, \quad (\text{see [10, 11, 13]}).$$

When $k = 1$, we have

$$(1.14) \quad \begin{aligned} \sum_{n=0}^{\infty} B_n^{(1)}(x) \frac{t^n}{n!} &= \frac{t}{1 - e^{-t}} e^{xt} = \frac{t}{e^t - 1} e^{(x+1)t} \\ &= \sum_{n=0}^{\infty} B_n(x+1) \frac{t^n}{n!}. \end{aligned}$$

By (1.14), we easily get

$$B_n^{(1)}(x) = B_n(x+1), \quad (n \geq 0).$$

Let $x = 0$. Then $B_n^{(k)} = B_n^{(k)}(0)$ are called the poly-Bernoulli numbers.

In this paper, we introduce the new fully degenerate poly-Bernoulli numbers and polynomials and investigate some properties of these polynomials and numbers. From our investigation, we derive some identities for the fully degenerate poly-Bernoulli numbers and polynomials.

2. FULLY DEGENERATE POLY-BERNOULLI POLYNOMIALS

For $k \in \mathbb{Z}$, we define the fully degenerate poly-Bernoulli polynomials which are given by the generating function

$$(2.1) \quad \frac{\text{Li}_k \left(1 - (1 + \lambda t)^{-\frac{1}{\lambda}} \right)}{1 - (1 + \lambda t)^{-\frac{1}{\lambda}}} (1 + \lambda t)^{\frac{x}{\lambda}} = \sum_{n=0}^{\infty} \beta_{n,\lambda}^{(k)}(x) \frac{t^n}{n!}.$$

When $x = 0$, $\beta_{n,\lambda}^{(k)} = \beta_{n,\lambda}^{(k)}(0)$ are called the fully degenerate poly-Bernoulli numbers.

From (1.13) and (2.1), we have

$$(2.2) \quad \begin{aligned} \frac{\text{Li}_k(1 - e^{-t})}{1 - e^{-t}} e^{xt} &= \lim_{\lambda \rightarrow 0} \frac{\text{Li}_k \left(1 - (1 + \lambda t)^{-\frac{1}{\lambda}} \right)}{1 - (1 + \lambda t)^{-\frac{1}{\lambda}}} (1 + \lambda t)^{\frac{x}{\lambda}} \\ &= \sum_{n=0}^{\infty} \lim_{\lambda \rightarrow 0} \beta_{n,\lambda}^{(k)}(x) \frac{t^n}{n!}. \end{aligned}$$

Thus, we get

$$(2.3) \quad \lim_{\lambda \rightarrow 0} \beta_{n,\lambda}^{(k)}(x) = B_n^{(k)}(x), \quad (n \geq 0).$$

By (2.1), we get

$$(2.4) \quad \begin{aligned} &\sum_{n=0}^{\infty} \beta_{n,\lambda}^{(k)}(x) \frac{t^n}{n!} \\ &= \frac{\text{Li}_k \left(1 - (1 + \lambda t)^{-\frac{1}{\lambda}} \right)}{1 - (1 + \lambda t)^{-\frac{1}{\lambda}}} (1 + \lambda t)^{\frac{x}{\lambda}} \\ &= \sum_{n=0}^{\infty} \left(\sum_{l=0}^n \binom{n}{l} \beta_{l,\lambda}^{(k)} \left(\frac{x}{\lambda} \right) \lambda^{n-l} \right) \frac{t^n}{n!}. \end{aligned}$$

Thus, from (2.4), we have

$$(2.5) \quad \beta_{n,\lambda}^{(k)}(x + y) = \sum_{l=0}^n \binom{n}{l} \left(\frac{y}{\lambda} \right)_{n-l} \lambda^{n-l} \beta_{l,\lambda}^{(k)}(x), \quad (n \geq 0),$$

and

$$\beta_{n,\lambda}^{(k)}(x) = \sum_{l=0}^n \binom{n}{l} \left(\frac{x}{\lambda} \right)_{n-l} \lambda^{n-l} \beta_{l,\lambda}^{(k)}.$$

Therefore, by (2.3) and (2.5), we obtain the following theorem.

Theorem 1. For $k \in \mathbb{Z}$, $n \geq 0$, we have

$$(2.6) \quad \beta_{n,\lambda}^{(k)}(x+y) = \sum_{l=0}^n \binom{n}{l} \left(\frac{y}{\lambda}\right)_{n-l} \lambda^{n-l} \beta_{l,\lambda}^{(k)}(x), \quad (n \geq 0),$$

and

$$\lim_{\lambda \rightarrow 0} \beta_{n,\lambda}^{(k)}(x) = B_n^{(k)}(x),$$

where $(x)_n = x(x-1) \cdots (x-n+1) = \sum_{l=0}^n S_1(n, l) x^l$.

From (2.1), we can derive the following equation:

$$(2.7) \quad \sum_{n=0}^{\infty} \beta_{n,\lambda}^{(k)}(x) \frac{t^n}{n!} = \frac{\text{Li}_k \left(1 - (1 + \lambda t)^{-\frac{1}{\lambda}} \right)}{(1 + \lambda t)^{\frac{1}{\lambda}} - 1} (1 + \lambda t)^{\frac{x+1}{\lambda}}.$$

Thus, by (2.7), we get

$$(2.8) \quad \begin{aligned} & \sum_{n=0}^{\infty} \left\{ \beta_{n,\lambda}^{(k)} - \beta_{n,\lambda}^{(k)}(-1) \right\} \frac{t^n}{n!} \\ &= \text{Li}_k \left(1 - (1 + \lambda t)^{-\frac{1}{\lambda}} \right) \\ &= \sum_{m=1}^{\infty} \frac{\left(1 - (1 + \lambda t)^{-\frac{1}{\lambda}} \right)^m}{m^k} \\ &= \sum_{m=0}^{\infty} \frac{(-1)^{m+1}}{(m+1)^k} \left(e^{-\frac{1}{\lambda} \log(1+\lambda t)} - 1 \right)^{m+1} \\ &= \sum_{m=0}^{\infty} \frac{(-1)^{m+1}}{(m+1)^k} (m+1)! \sum_{l=m+1}^{\infty} S_2(l, m+1) (-1)^l \lambda^{-l} \frac{(\log(1+\lambda t))^l}{l!} \\ &= \sum_{l=1}^{\infty} \sum_{m=0}^{l-1} \frac{(-1)^{m+1}}{(m+1)^k} (m+1)! S_2(l, m+1) (-1)^l \lambda^{-l} \sum_{n=l}^{\infty} S_1(n, l) \lambda^n \frac{t^n}{n!} \\ &= \sum_{n=1}^{\infty} \left(\sum_{l=1}^n \sum_{m=0}^{l-1} \frac{m! (m+1) (-1)^{l-m-1} \lambda^{n-l} S_2(l, m+1) S_1(n, l)}{(m+1)^k} \right) \frac{t^n}{n!}, \end{aligned}$$

where $S_2(n, l)$ and $S_1(n, l)$ are the Stirling numbers of the second kind and of the first kind, respectively.

Therefore, by (2.8), we obtain the following theorem.

Theorem 2. For $k \in \mathbb{Z}$, $n \geq 1$, we have

$$\beta_{n,\lambda}^{(k)} - \beta_{n,\lambda}^{(k)}(-1) = \sum_{l=1}^n \sum_{m=0}^{l-1} \frac{m! (-1)^{l-m-1} \lambda^{n-l} S_2(l, m+1) S_1(n, l)}{(m+1)^{k-1}}.$$

From (1.12), we can easily derive the following equation:

$$(2.9) \quad \text{Li}'_k(t) = \frac{d}{dt} \text{Li}_k(t) = \frac{1}{t} \text{Li}_{k-1}(t).$$

Thus, by (2.9), the generating function of the fully degenerate poly-Bernoulli numbers is also written in terms of the following iterated integral:

$$\begin{aligned}
 (2.10) \quad & \frac{(1+\lambda t)^{\frac{1}{\lambda}}}{(1+\lambda t)^{\frac{1}{\lambda}} - 1} \int_0^t \frac{1}{\left((1+\lambda t)^{\frac{1}{\lambda}} - 1\right)(1+\lambda t)} \\
 & \times \int_0^t \frac{1}{\left((1+\lambda t)^{\frac{1}{\lambda}} - 1\right)(1+\lambda t)} \cdots \int_0^t \frac{\log(1+\lambda t)^{\frac{1}{\lambda}}}{\left((1+\lambda t)^{\frac{1}{\lambda}} - 1\right)(1+\lambda t)} \underbrace{dt \cdots dt}_{k-1 \text{ times}} \\
 & = \sum_{n=0}^{\infty} \beta_{n,\lambda}^{(k)} \frac{t^n}{n!}.
 \end{aligned}$$

For $k = 2$, we have

$$\begin{aligned}
 (2.11) \quad & \sum_{n=0}^{\infty} \beta_{n,\lambda}^{(2)} \frac{t^n}{n!} \\
 & = \frac{(1+\lambda t)^{\frac{1}{\lambda}}}{(1+\lambda t)^{\frac{1}{\lambda}} - 1} \int_0^t \frac{\log(1+\lambda t)^{\frac{1}{\lambda}}}{(1+\lambda t)^{\frac{1}{\lambda}} - 1} (1+\lambda t)^{-\frac{1}{\lambda}} dt \\
 & = \frac{(1+\lambda t)^{\frac{1}{\lambda}}}{(1+\lambda t)^{\frac{1}{\lambda}} - 1} \left(\sum_{m=0}^{\infty} b_{m,\lambda}(-\lambda) \frac{1}{m!} \int_0^t t^m dt \right) \\
 & = \left(\frac{t}{(1+\lambda t)^{\frac{1}{\lambda}} - 1} (1+\lambda t)^{\frac{1}{\lambda}} \right) \left(\sum_{m=0}^{\infty} \frac{b_{m,\lambda}(-\lambda)}{(m+1)} \frac{t^m}{m!} \right) \\
 & = \sum_{n=0}^{\infty} \left(\sum_{l=0}^n \binom{n}{l} \beta_{l,\lambda}(1) \frac{b_{n-l,\lambda}(-\lambda)}{n-l+1} \right) \frac{t^n}{n!}.
 \end{aligned}$$

Therefore, by (2.11), we obtain the following theorem.

Theorem 3. For $n \geq 0$, we have

$$\beta_{n,\lambda}^{(2)} = \sum_{l=0}^n \binom{n}{l} \beta_{l,\lambda}(1) \frac{b_{n-l,\lambda}(-\lambda)}{n-l+1}.$$

Note that

$$B_n^{(2)} = \lim_{\lambda \rightarrow 0} \beta_{n,\lambda}^{(2)} = \sum_{l=0}^n \binom{n}{l} B_l(1) \frac{B_{n-l}}{n-l+1}.$$

From (2.1), we have

$$\begin{aligned}
 (2.12) \quad & \sum_{n=0}^{\infty} \beta_{n,\lambda}^{(k)} \frac{t^n}{n!} \\
 & = \frac{\text{Li}_k \left(1 - (1+\lambda t)^{-\frac{1}{\lambda}} \right)}{1 - (1+\lambda t)^{-\frac{1}{\lambda}}} \\
 & = \sum_{m=0}^{\infty} \frac{1}{(m+1)^k} \left(1 - (1+\lambda t)^{-\frac{1}{\lambda}} \right)^m \\
 & = \sum_{m=0}^{\infty} \frac{(-1)^m}{(m+1)^k} \left(e^{-\frac{1}{\lambda} \log(1+\lambda t)} - 1 \right)^m
 \end{aligned}$$

$$\begin{aligned}
&= \sum_{m=0}^{\infty} \frac{(-1)^m}{(m+1)^k} m! \sum_{l=m}^{\infty} S_2(l, m) \left(-\frac{1}{\lambda}\right)^l \frac{(\log(1+\lambda t))^l}{l!} \\
&= \sum_{l=0}^{\infty} \left(\sum_{m=0}^l \frac{(-1)^{m+l} m!}{(m+1)^k} S_2(l, m) \lambda^{-l} \right) \frac{1}{l!} (\log(1+\lambda t))^l \\
&= \sum_{l=0}^{\infty} \left(\sum_{m=0}^l \frac{(-1)^{m+l} m!}{(m+1)^k} S_2(l, m) \lambda^{-l} \right) \sum_{n=l}^{\infty} S_1(n, l) \lambda^n \frac{t^n}{n!} \\
&= \sum_{n=0}^{\infty} \left\{ \sum_{l=0}^n \sum_{m=0}^l \frac{(-1)^{m+l} m!}{(m+1)^k} S_2(l, m) S_1(n, l) \lambda^{n-l} \right\} \frac{t^n}{n!}.
\end{aligned}$$

Therefore, by (2.12), we obtain the following theorem.

Theorem 4. *For $n \geq 0$, we have*

$$\beta_{n,\lambda}^{(k)} = \sum_{l=0}^n \sum_{m=0}^l \frac{(-1)^{m+l} m!}{(m+1)^k} S_2(l, m) S_1(n, l) \lambda^{n-l}.$$

Note that

$$B_n^{(k)} = \lim_{\lambda \rightarrow 0} \beta_{n,\lambda}^{(k)} = \sum_{m=0}^n \frac{(-1)^{m+n} m!}{(m+1)^k} S_2(n, m).$$

From (2.9), we have

$$\begin{aligned}
(2.13) \quad & \frac{d}{dt} \text{Li}_k \left(1 - (1 + \lambda t)^{-\frac{1}{\lambda}} \right) \\
&= \frac{1}{1 - (1 + \lambda t)^{-\frac{1}{\lambda}}} (1 + \lambda t)^{-\frac{1}{\lambda}-1} \text{Li}_{k-1} \left(1 - (1 + \lambda t)^{-\frac{1}{\lambda}} \right) \\
&= (1 + \lambda t)^{-\frac{1}{\lambda}-1} \sum_{n=0}^{\infty} \beta_{n,\lambda}^{(k-1)} \frac{t^n}{n!}.
\end{aligned}$$

On the other hand,

$$\begin{aligned}
(2.14) \quad & \frac{d}{dt} \left(\text{Li}_k \left(1 - (1 + \lambda t)^{-\frac{1}{\lambda}} \right) \right) \\
&= \frac{d}{dt} \left(\left(1 - (1 + \lambda t)^{-\frac{1}{\lambda}} \right) \frac{1}{1 - (1 + \lambda t)^{-\frac{1}{\lambda}}} \text{Li}_k \left(1 - (1 + \lambda t)^{-\frac{1}{\lambda}} \right) \right) \\
&= (1 + \lambda t)^{-\frac{1}{\lambda}-1} \frac{1}{1 - (1 + \lambda t)^{-\frac{1}{\lambda}}} \text{Li}_k \left(1 - (1 + \lambda t)^{-\frac{1}{\lambda}} \right) \\
&\quad + \left(1 - (1 + \lambda t)^{-\frac{1}{\lambda}} \right) \frac{d}{dt} \left(\sum_{n=0}^{\infty} \beta_{n,\lambda}^{(k)} \frac{t^n}{n!} \right) \\
&= (1 + \lambda t)^{-\frac{1}{\lambda}-1} \sum_{n=0}^{\infty} \beta_{n,\lambda}^{(k)} \frac{t^n}{n!} \\
&\quad + \left(1 - (1 + \lambda t)^{-\frac{1}{\lambda}} \right) \sum_{n=1}^{\infty} \beta_{n,\lambda}^{(k)} \frac{t^{n-1}}{(n-1)!}.
\end{aligned}$$

By (2.13) and (2.14), we get

$$\begin{aligned}
(2.15) \quad & \sum_{n=0}^{\infty} \beta_{n,\lambda}^{(k-1)} \frac{t^n}{n!} \\
&= \sum_{n=0}^{\infty} \beta_{n,\lambda}^{(k)} \frac{t^n}{n!} + (1+\lambda t) \left((1+\lambda t)^{\frac{1}{\lambda}} - 1 \right) \sum_{n=1}^{\infty} \beta_{n,\lambda}^{(k)} \frac{t^{n-1}}{(n-1)!} \\
&= \sum_{n=0}^{\infty} \beta_{n,\lambda}^{(k)} \frac{t^n}{n!} + \left((1+\lambda t)^{\frac{1}{\lambda}} - 1 \right) \sum_{m=0}^{\infty} \beta_{m+1,\lambda}^{(k)} \frac{t^m}{m!} \\
&\quad + \lambda \left((1+\lambda t)^{\frac{1}{\lambda}} - 1 \right) \sum_{m=0}^{\infty} \beta_{m,\lambda}^{(k)} m \frac{t^m}{m!} \\
&= \sum_{n=0}^{\infty} \beta_{n,\lambda}^{(k)} \frac{t^n}{n!} + \left(\sum_{l=1}^{\infty} \left(\frac{1}{\lambda} \right)_l \lambda^l \frac{t^l}{l!} \right) \left(\sum_{m=0}^{\infty} \beta_{m+1,\lambda}^{(k)} \frac{t^m}{m!} \right) \\
&\quad + \lambda \left(\sum_{l=1}^{\infty} \left(\frac{1}{\lambda} \right)_l \lambda^l \frac{t^l}{l!} \right) \left(\sum_{m=0}^{\infty} \beta_{m,\lambda}^{(k)} m \frac{t^m}{m!} \right) \\
&= \sum_{n=0}^{\infty} \beta_{n,\lambda}^{(k)} \frac{t^n}{n!} + \sum_{n=1}^{\infty} \left(\sum_{m=0}^{n-1} \left(\frac{1}{\lambda} \right)_{n-m} \lambda^{n-m} \frac{n!}{(n-m)!m!} \beta_{m+1,\lambda}^{(k)} \right) \frac{t^n}{n!} \\
&\quad + \lambda \sum_{n=1}^{\infty} \left(\sum_{m=0}^{n-1} \left(\frac{1}{\lambda} \right)_{n-m} \lambda^{n-m} \frac{m \cdot n!}{(n-m)!m!} \beta_{m,\lambda}^{(k)} \right) \frac{t^n}{n!} \\
&= \sum_{n=0}^{\infty} \beta_{n,\lambda}^{(k)} \frac{t^n}{n!} + \sum_{n=1}^{\infty} \left(\sum_{m=0}^{n-1} \binom{n}{m} \left(\frac{1}{\lambda} \right)_{n-m} \lambda^{n-m} \beta_{m+1,\lambda}^{(k)} \right) \frac{t^n}{n!} \\
&\quad + \lambda \sum_{n=1}^{\infty} \left(\sum_{m=0}^{n-1} \binom{n}{m} \left(\frac{1}{\lambda} \right)_{n-m} \lambda^{n-m} m \beta_{m,\lambda}^{(k)} \right) \frac{t^n}{n!},
\end{aligned}$$

where

$$\left(\frac{1}{\lambda} \right)_n = \left(\frac{1}{\lambda} \right) \left(\frac{1}{\lambda} - 1 \right) \cdots \left(\frac{1}{\lambda} - n + 1 \right) = \sum_{l=0}^n S_1(n, l) \lambda^{-l}, \quad (n \geq 0).$$

Thus, by (2.15), we have

$$\begin{aligned}
(2.16) \quad & \beta_{n,\lambda}^{(k-1)} = \beta_{n,\lambda}^{(k)} + \sum_{m=0}^{n-1} \binom{n}{m} \left(\frac{1}{\lambda} \right)_{n-m} \lambda^{n-m} \beta_{m+1,\lambda}^{(k)} \\
&\quad + \lambda \sum_{m=0}^{n-1} \binom{n}{m} \left(\frac{1}{\lambda} \right)_{n-m} \lambda^{n-m} m \beta_{m,\lambda}^{(k)} \\
&= (n+1) \beta_{n,\lambda}^{(k)} + \sum_{m=1}^{n-1} \binom{n}{m-1} \left(\frac{1}{\lambda} \right)_{n-m+1} \lambda^{n-m+1} \beta_{m,\lambda}^{(k)} \\
&\quad + \lambda \sum_{m=0}^{n-1} \binom{n}{m} \left(\frac{1}{\lambda} \right)_{n-m} \lambda^{n-m} m \beta_{m,\lambda}^{(k)}, \quad (n \geq 1).
\end{aligned}$$

Therefore, by (2.16), we obtain the following theorem.

Theorem 5. *For $n \geq 1$, we have*

$$\beta_{n,\lambda}^{(k)} = \frac{1}{n+1} \left\{ \beta_{n,\lambda}^{(k-1)} - \sum_{m=1}^{n-1} \binom{n}{m-1} \beta_{m,\lambda}^{(k)} \left(\frac{1}{\lambda} \right)_{n-m+1} \lambda^{n-m+1} \right. \\ \left. - \lambda \sum_{m=0}^{n-1} \binom{n}{m} \left(\frac{1}{\lambda} \right)_{n-m} \lambda^{n-m} m \beta_{m,\lambda}^{(k)} \right\}.$$

Note that

$$B_n^{(k)} = \lim_{\lambda \rightarrow 0} \beta_{n,\lambda}^{(k)} = \frac{1}{n+1} \left\{ B_n^{(k-1)} - \sum_{m=1}^{n-1} \binom{n}{m-1} B_m^{(k)} \right\}.$$

Now, we observe that

$$(2.17) \quad \sum_{n=0}^{\infty} \left(1 - (1 + \lambda t)^{-\frac{1}{\lambda}} \right)^n (n+1)^{-k} \\ = \sum_{n=1}^{\infty} \frac{\left(1 - (1 + \lambda t)^{-\frac{1}{\lambda}} \right)^n}{n^k} \frac{1}{1 - (1 + \lambda t)^{-\frac{1}{\lambda}}} \\ = \frac{1}{1 - (1 + \lambda t)^{-\frac{1}{\lambda}}} \text{Li}_k \left(1 - (1 + \lambda t)^{-\frac{1}{\lambda}} \right) \\ = \sum_{n=0}^{\infty} \beta_{n,\lambda}^{(k)} \frac{t^n}{n!}.$$

By (2.17), we get

$$(2.18) \quad \sum_{k=0}^{\infty} \left(\sum_{n=0}^{\infty} \beta_{n,\lambda}^{(-k)} \frac{x^n}{n!} \right) \frac{y^k}{k!} \\ = \sum_{k=0}^{\infty} \left(\sum_{m=0}^{\infty} \left(1 - (1 + \lambda t)^{-\frac{1}{\lambda}} \right)^m (m+1)^k \right) \frac{y^k}{k!} \\ = \sum_{m=0}^{\infty} \left(1 - (1 + \lambda x)^{-\frac{1}{\lambda}} \right)^m \sum_{k=0}^{\infty} (m+1)^k \frac{y^k}{k!} \\ = \sum_{m=0}^{\infty} \left(1 - (1 + \lambda x)^{-\frac{1}{\lambda}} \right)^m e^{(m+1)y} \\ = \sum_{j=0}^{\infty} (-1)^j \left(e^{-\frac{1}{\lambda} \log(1+\lambda x)} - 1 \right)^j e^{(j+1)y} \\ = \sum_{j=0}^{\infty} (-1)^j j! \sum_{m=j}^{\infty} S_2(m, j) (-1)^m \lambda^{-m} \frac{(\log(1+\lambda x))^m}{m!} e^{(j+1)y} \\ = \sum_{m=0}^{\infty} \sum_{j=0}^m (-1)^{j+m} j! S_2(m, j) \lambda^{-m} \sum_{n=m}^{\infty} S_1(n, m) \lambda^n \frac{x^n}{n!} e^{(j+1)y}$$

$$\begin{aligned}
&= \sum_{n=0}^{\infty} \left(\sum_{m=0}^n \sum_{j=0}^m (-1)^{j+m} j! S_2(m, j) \lambda^{n-m} S_1(n, m) e^{(j+1)y} \right) \frac{x^n}{n!} \\
&= \sum_{k=0}^{\infty} \sum_{n=0}^{\infty} \left(\sum_{m=0}^n \sum_{j=0}^m (-1)^{j+m} j! \lambda^{n-m} S_2(m, j) S_1(n, m) (j+1)^k \right) \frac{x^n}{n!} \frac{y^k}{k!}.
\end{aligned}$$

Therefore, by (2.18), we obtain the following theorem.

Theorem 6. For $k \in \mathbb{Z}$ and $n \geq 0$, we have

$$\beta_{n,\lambda}^{(-k)} = \sum_{m=0}^n \sum_{j=0}^m (-1)^{j+m} j! \lambda^{n-m} (j+1)^k S_2(m, j) S_1(n, m).$$

Note that

$$B_n^{(-k)} = \lim_{\lambda \rightarrow 0} \beta_{n,\lambda}^{(-k)} = \sum_{j=0}^n (-1)^{j+n} j! (j+1)^k S_2(n, j).$$

From Theorem 1, we have

$$\begin{aligned}
\frac{d}{dx} \beta_{n,\lambda}^{(k)}(x) &= \sum_{l=0}^n \binom{n}{l} \beta_{l,\lambda}^{(k)} \frac{d}{dx} \left(\prod_{i=0}^{n-l-1} (x - i\lambda) \right) \\
&= \sum_{l=0}^n \binom{n}{l} \beta_{l,\lambda}^{(k)} \sum_{j=0}^{n-l-1} \frac{1}{(x - j\lambda)} \prod_{i=0}^{n-l-1} (x - i\lambda).
\end{aligned}$$

The generalized falling factorial $(x | \lambda)_n$ is given by

$$(2.19) \quad (x | \lambda)_n = x(x - \lambda)(x - 2\lambda) \cdots (x - (n-1)\lambda), \quad (n \geq 0).$$

As is well known, the Bernoulli numbers of the second kind are defined by the generating function

$$(2.20) \quad \frac{t}{\log(1+t)} = \sum_{n=0}^{\infty} b_n \frac{t^n}{n!}, \quad (\text{see [20]}).$$

We observe that

$$\begin{aligned}
(2.21) \quad \int_0^1 (1 + \lambda t)^{\frac{x}{\lambda}} dx &= \sum_{n=0}^{\infty} \lambda^n \int_0^1 \left(\frac{x}{\lambda} \right)_n dx \frac{t^n}{n!} \\
&= \sum_{n=0}^{\infty} \int_0^1 (x | \lambda)_n dx \frac{t^n}{n!}.
\end{aligned}$$

On the other hand,

$$\begin{aligned}
(2.22) \quad \int_0^1 (1 + \lambda t)^{\frac{x}{\lambda}} dx &= \frac{\lambda}{\log(1 + \lambda t)} \left((1 + \lambda t)^{\frac{1}{\lambda}} - 1 \right) \\
&= \frac{\lambda t}{\log(1 + \lambda t)} \left(\frac{(1 + \lambda t)^{\frac{1}{\lambda}} - 1}{t} \right)
\end{aligned}$$

$$\begin{aligned}
&= \left(\sum_{m=0}^{\infty} b_m \lambda^m \frac{t^m}{m!} \right) \left(\sum_{l=0}^{\infty} \frac{(1|\lambda)_{l+1}}{l+1} \frac{t^l}{l!} \right) \\
&= \sum_{n=0}^{\infty} \left(\sum_{l=0}^n \frac{(1|\lambda)_{l+1}}{l+1} \lambda^{n-l} b_{n-l} \binom{n}{l} \right) \frac{t^n}{n!}.
\end{aligned}$$

From (2.21) and (2.22), we have

$$(2.23) \quad \int_0^1 (x|\lambda)_n dx = \sum_{l=0}^n \binom{n}{l} \lambda^{n-l} b_{n-l} \frac{(1|\lambda)_{l+1}}{l+1}, \quad (n \geq 0).$$

By Theorem 1, we get

$$\begin{aligned}
\int_0^1 \beta_{n,\lambda}^{(k)}(x) dx &= \sum_{l=0}^n \binom{n}{l} \beta_{l,\lambda}^{(k)} \int_0^1 \left(\frac{x}{\lambda}\right)_{n-l} \lambda^{n-l} dx \\
&= \sum_{l=0}^n \binom{n}{l} \beta_{n-l,\lambda}^{(k)} \int_0^1 (x|\lambda)_l dx \\
&= \sum_{l=0}^n \left(\sum_{m=0}^l \binom{l}{m} \lambda^{l-m} b_{l-m} \frac{(1|\lambda)_{m+1}}{m+1} \right) \binom{n}{l} \beta_{n-l,\lambda}^{(k)}.
\end{aligned}$$

3. FURTHER REMARKS

Let $\mathbb{C}$ be complex number field and let $\mathcal{F}$ be the set of all formal power series in the variable t over $\mathbb{C}$ with

$$(3.1) \quad \mathcal{F} = \left\{ f(t) = \sum_{k=0}^{\infty} a_k \frac{t^k}{k!} \mid a_k \in \mathbb{C} \right\}.$$

Let $\mathbb{P}$ be the algebra of polynomials in a single variable x over $\mathbb{C}$ and let $\mathbb{P}^*$ be the vector space of all linear functionals on $\mathbb{P}$. The action of linear functional $L \in \mathbb{P}^*$ on a polynomial $p(x)$ is denoted by $\langle L|p(x) \rangle$, and linearly extended as

$$\langle cL + c'L'|p(x) \rangle = c \langle L|p(x) \rangle + c' \langle L'|p(x) \rangle,$$

where $c, c' \in \mathbb{C}$.

For $f(t) = \sum_{k=0}^{\infty} a_k \frac{t^k}{k!}$, we define a linear functional on $\mathbb{P}$ by setting

$$(3.2) \quad \langle f(t)|x^n \rangle = a_n \quad \text{for all } n \geq 0.$$

Thus, by (3.2), we get

$$(3.3) \quad \langle t^k|x^n \rangle = n! \delta_{n,k}, \quad (n, k \geq 0), \quad (\text{see [6, 17, 20]}).$$

For $f_L(t) = \sum_{k=0}^{\infty} \langle L|x^k \rangle \frac{t^k}{k!}$, by (3.3), we get $\langle f_L(t)|x^n \rangle = \langle L|x^n \rangle$. In addition, the mapping $L \mapsto f_L(t)$ is a vector space isomorphism from $\mathbb{P}^*$ onto $\mathcal{F}$. Henceforth, $\mathcal{F}$ denotes both the algebra of the formal power series in t and the vector space of all linear functionals on $\mathbb{P}$ and so an element $f(t)$ of $\mathcal{F}$ can be regarded as both a formal power series and a linear functional. We refer to $\mathcal{F}$ umbral algebra. The umbral calculus is the study of umbral algebra (see [9, 16, 20]). The order $o(f(t))$ of the non-zero power series $f(t)$ is the smallest integer k for which the coefficient of t^k does not vanish.

If $o(f(t)) = 1$ (respectively, $o(f(t)) = 0$), then $f(t)$ is called a delta (respectively, an invertible) series (see [20]). For $o(f(t)) = 1$ and $o(g(t)) = 0$, there exists

a unique sequence $s_n(x)$ of polynomials such that $\langle g(t)f(t)^k | s_n(x) \rangle = n! \delta_{n,k}$, ($n, k \geq 0$).

The sequence $s_n(x)$ is called the Sheffer sequence for $(g(t), f(t))$, and we write $s_n(x) \sim (g(t), f(t))$ (see [20]).

Let $f(t) \in \mathcal{F}$ and $p(x) \in \mathbb{P}$. Then, by (3.3), we get

$$(3.4) \quad f(t) = \sum_{k=0}^{\infty} \langle f(t) | x^k \rangle \frac{t^k}{k!}, \quad p(x) = \sum_{k=0}^{\infty} \langle t^k | p(x) \rangle \frac{x^k}{k!}.$$

From (3.4), we have

$$(3.5) \quad p^{(k)}(0) = \langle t^k | p(x) \rangle = \langle 1 | p^{(k)}(x) \rangle,$$

where $p^k(x) = \frac{d^k}{dx^k} p(x)$, (see [12, 15, 20]).

By (3.5), we easily get

$$(3.6) \quad t^k p(x) = p^k(x), \quad e^{yt} p(x) = p(x+y), \quad \text{and} \quad \langle e^{yt} | p(x) \rangle = p(y).$$

From (3.6), we have

$$\frac{e^{yt} - 1}{t} p(x) = \int_x^{x+y} p(u) du, \quad \langle e^{yt} - 1 | p(x) \rangle = p(y) - p(0).$$

Let $f(t)$ be the linear functional such that

$$(3.7) \quad \langle f(t) | p(x) \rangle = \int_0^y p(u) du,$$

for all polynomials $p(x)$. Then it can be determined by (3.4) to be

$$(3.8) \quad \begin{aligned} f(t) &= \sum_{k=0}^{\infty} \frac{\langle f(t) | x^k \rangle}{k!} t^k \\ &= \sum_{k=0}^{\infty} \frac{y^{k+1}}{(k+1)!} t^k \\ &= \frac{1}{t} (e^{yt} - 1). \end{aligned}$$

Thus, for $p(x) \in \mathbb{P}$, we have

$$(3.9) \quad \left\langle \frac{e^{yt} - 1}{t} \middle| p(x) \right\rangle = \int_0^y p(u) du.$$

It is known that

$$(3.10) \quad s_n(x) \sim (g(t), f(t)) \iff \frac{1}{g(\bar{f}(t))} e^{x\bar{f}(t)} = \sum_{k=0}^{\infty} s_k(x) \frac{t^k}{k!}, \quad (x \in \mathbb{C})$$

where $\bar{f}(t)$ is the compositional inverse of $f(t)$ such that $f(\bar{f}(t)) = \bar{f}(f(t)) = t$ (see [12, 20]).

From (2.1), we note that

$$(3.11) \quad \beta_{n,\lambda}^{(k)}(x) \sim \left(\frac{1 - e^{-t}}{\text{Li}_k(1 - e^{-t})}, \frac{1}{\lambda} (e^{\lambda t} - 1) \right).$$

That is,

$$\sum_{n=0}^{\infty} \beta_{n,\lambda}^{(k)}(x) \frac{t^n}{n!} = \frac{\text{Li}_k \left(1 - (1 + \lambda t)^{-\frac{1}{\lambda}} \right)}{1 - (1 + \lambda t)^{-\frac{1}{\lambda}}} (1 + \lambda t)^{\frac{x}{\lambda}}.$$

Thus, by (3.11),

$$(3.12) \quad \frac{1}{\lambda} (e^{\lambda t} - 1) \beta_{n,\lambda}^{(k)}(x) = n \beta_{n-1,\lambda}^{(k)}(x).$$

On the other hand,

$$(3.13) \quad (e^{\lambda t} - 1) \beta_{n,\lambda}^{(k)}(x) = \beta_{n,\lambda}^{(k)}(x + \lambda) - \beta_{n,\lambda}^{(k)}(x).$$

Therefore, by (3.12) and (3.13), we obtain the following theorem.

Theorem 7. *For $n \in \mathbb{N}$, we have*

$$\lambda \beta_{n-1,\lambda}^{(k)}(x) = \frac{1}{n} \left\{ \beta_{n,\lambda}^{(k)}(x + \lambda) - \beta_{n,\lambda}^{(k)}(x) \right\}.$$

By (3.9), we get

$$(3.14) \quad \frac{e^{yt} - 1}{t} \beta_{n,\lambda}^{(k)}(x) = \int_x^{x+y} \beta_{n,\lambda}^{(k)}(u) du.$$

From (3.14), we have

$$(3.15) \quad \left\langle \frac{e^{yt} - 1}{t} \middle| \beta_{n,\lambda}^{(k)}(x) \right\rangle = \int_0^y \beta_{n,\lambda}^{(k)}(u) du.$$

Thus, by (3.15), we get

$$(3.16) \quad \begin{aligned} & \left\langle \frac{e^t - 1}{t} \middle| \beta_{n,\lambda}^{(k)}(x) \right\rangle \\ &= \int_0^1 \beta_{n,\lambda}^{(k)}(u) du \\ &= \sum_{l=0}^n \sum_{m=0}^l \binom{l}{m} \binom{n}{l} \lambda^{l-m} b_{l-m} \beta_{n-l,\lambda}^{(k)} \frac{(1 | \lambda)_{m+1}}{m+1}. \end{aligned}$$

Therefore, by (3.16), we obtain the following theorem.,

Theorem 8. *For $n \geq 0$, we have*

$$\left\langle \frac{e^t - 1}{t} \middle| \beta_{n,\lambda}^{(k)}(x) \right\rangle = \sum_{l=0}^n \sum_{m=0}^l \binom{l}{m} \binom{n}{l} \lambda^{l-m} b_{l-m} \beta_{n-l,\lambda}^{(k)} \frac{(1 | \lambda)_{m+1}}{m+1}.$$

Note that

$$\begin{aligned} & \left\langle \frac{e^t - 1}{t} \middle| B_n^{(k)}(x) \right\rangle \\ &= \lim_{\lambda \rightarrow 0} \left\langle \frac{e^t - 1}{t} \middle| \beta_{n,\lambda}^{(k)}(x) \right\rangle \\ &= \lim_{\lambda \rightarrow 0} \int_0^1 \beta_{n,\lambda}^{(k)}(u) du \\ &= \sum_{l=0}^n \binom{n}{l} B_{n-l}^{(k)} \frac{1}{l+1} \end{aligned}$$

Let

$$\mathbb{P}_n = \{p(x) \in \mathbb{C}[x] \mid \deg p(x) \leq n\}, \quad (n \geq 0).$$

For $p(x) \in \mathbb{P}_n$ with $p(x) = \sum_{m=0}^n a_m \beta_{m,\lambda}^{(k)}(x)$, we have

$$(3.17) \quad \begin{aligned} & \left\langle \frac{1-e^{-t}}{\text{Li}_k(1-e^{-t})} \left(\frac{1}{\lambda} (e^{\lambda t} - 1) \right)^m \middle| p(x) \right\rangle \\ &= \sum_{l=0}^n a_l \left\langle \frac{1-e^{-t}}{\text{Li}_k(1-e^{-t})} \left(\frac{1}{\lambda} (e^{\lambda t} - 1) \right)^m \middle| \beta_{l,\lambda}^{(k)}(x) \right\rangle \end{aligned}$$

From (3.11), we note that

$$(3.18) \quad \left\langle \frac{1-e^{-t}}{\text{Li}_k(1-e^{-t})} \left(\frac{1}{\lambda} (e^{\lambda t} - 1) \right)^m \middle| \beta_{l,\lambda}^{(k)}(x) \right\rangle = l! \delta_{l,m}.$$

By (3.17) and (3.18), we get

$$(3.19) \quad a_m = \frac{1}{m!} \left\langle \frac{1-e^{-t}}{\text{Li}_k(1-e^{-t})} \left(\frac{1}{\lambda} (e^{\lambda t} - 1) \right)^m \middle| p(x) \right\rangle, \quad (m \geq 0).$$

Therefore, by (3.19), we obtain the following theorem.

Theorem 9. For $p(x) \in \mathbb{P}_n$, we have

$$p(x) = \sum_{m=0}^n a_m \beta_{m,\lambda}^{(k)}(x),$$

where

$$a_m = \frac{1}{m!} \left\langle \frac{1-e^{-t}}{\text{Li}_k(1-e^{-t})} \left(\frac{1}{\lambda} (e^{\lambda t} - 1) \right)^m \middle| p(x) \right\rangle.$$

For example, let us take $p(x) = B_n^{(k)}(x)$ ($n \geq 0$). Then, by Theorem 9, we have

$$(3.20) \quad B_n^{(k)}(x) = \sum_{m=0}^n a_m \beta_{m,\lambda}^{(k)}(x),$$

where

$$(3.21) \quad \begin{aligned} a_m &= \frac{1}{m!} \left\langle \frac{1-e^{-t}}{\text{Li}_k(1-e^{-t})} \left(\frac{1}{\lambda} (e^{\lambda t} - 1) \right)^m \middle| B_n^{(k)}(x) \right\rangle \\ &= \frac{1}{m!} \left\langle \frac{1-e^{-t}}{\text{Li}_k(1-e^{-t})} \left(\frac{1}{\lambda} (e^{\lambda t} - 1) \right)^m \middle| \frac{\text{Li}_k(1-e^{-t})}{1-e^{-t}} x^n \right\rangle \\ &= \frac{\lambda^{-m}}{m!} \left\langle (e^{\lambda t} - 1)^m \middle| x^n \right\rangle = \lambda^{-m} \sum_{l=m}^{\infty} S_2(l, m) \frac{\lambda^l}{l!} \langle t^l | x^n \rangle \\ &= \lambda^{n-m} S_2(n, m). \end{aligned}$$

From (3.20) and (3.21), we have

$$(3.22) \quad B_n^{(k)}(x) = \sum_{m=0}^n \lambda^{n-m} S_2(n, m) \beta_{m,\lambda}^{(k)}(x).$$

REFERENCES

1. S. Araci, M. Acikgoz, and A. Kilicman, *Extended p -adic q -invariant integrals on $\mathbb{Z}_p$ associated with applications of umbral calculus*, Adv. Difference Equ. (2013), 2013:96, 14 pp.
2. A. Bayad, and T. Kim, *Results on values of Barnes polynomials*, Rocky Mountain J. Math. **43** (2013), 1857-1869.
3. J. Broedel, O. Schlotterer and S. Stieberger, *Polylogarithms, multiple zeta values and superstring amplitudes*, Fortschr. Phys. **61** (2013), no. 9, 812-870.
4. L. Carlitz, *Degenerate Stirling, Bernoulli and Eulerian numbers*, Utilitas Math. **15** (1979), 51-88.
5. D. Ding and J. Yang, *Some identities related to the Apostol-Euler and Apostol-Bernoulli polynomials*, Adv. Stud. Contemp. Math. (Kyungshang) **20** (2010), no. 1, 7-21.
6. H. Gangl, *Herbert Functional equations and ladders for polylogarithms*, Commun. Number Theory Phys. **7** (2013), no. 3, 397-410.
7. S. Gaboury, R. Tremblay and B.-J. Fugere, *Some explicit formulas for certain new classes of Bernoulli, Euler and Genocchi polynomials*, Proc. Jangjeon Math. Soc. **17** (2014), no. 1, 97-104.
8. Y. He and W. Zhang, *A convolution formula for Bernoulli polynomials*, Ars Combin. **108** (2013), 97-104.
9. H. Jolany, M. Aliabadi, R.B. Corcino and M.R. Darafsheh, *A note on multi poly-Euler numbers and Bernoulli polynomials*, General Mathematics **20** (2012), no. 2-3, 122-134.
10. M. Kaneko, *poly-Bernoulli numbers*, J Théor. Nombres Bordeaux **9** (1997), no. 1, 221-228.
11. D. S. Kim and T. Kim, *A note on poly-Bernoulli and higher-order poly-Bernoulli polynomials*, Russ. J. Math. Phys. **22** (2015), no. 1, 26-33.
12. ———, *Higher-order Cauchy of the first kind and poly-Cauchy of the first kind mixed type polynomials*, Adv. Stud. Contemp. Math. (Kyungshang) **23** (2013), no. 4, 621-636.
13. ———, *Higher-order Frobenius-Euler and poly-Bernoulli mixed-type polynomials*, Adv. Difference Equ. (2013), 2013:251, 13 pp.
14. ———, *Daehee polynomials with q -parameter*, Adv. Studies Theor. Phys. **8** (2014), no. 13, 561-569.
15. D. S. Kim, T. Kim, T. Mansour, and D. V. Dolgy, *On poly-Bernoulli polynomials of the second kind with umbral calculus viewpoint*, Adv. Difference Equ. (2015), 2015:27, 13 pp.
16. T. Kim, *Barnes' type multiple degenerate Bernoulli and Euler polynomials*, Appl. Math. Comput. (2015), 556-564.
17. T. Kim and T. Mansour, *Umbral calculus associated with Frobenius-type Eulerian polynomials*, Russ. J. Math. Phys. **21** (2014), no. 4, 484-493.
18. Q.-M. Luo, *Some recursion formula and relations for Bernoulli numbers and Euler numbers of higher order*, Adv. Stud. Contemp. Math. (Kyungshang) **10** (2005), no. 1, 63-70.
19. F. Qi, *An integral representation, complete monotonicity, and inequalities of Cauchy numbers of the second kind*, J. Number Theory **144** (2014), 244-255.
20. S. Roman, *The umbral calculus*, Pure and Applied Mathematics, vol. 111, Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York, 1984.

MR 741185 (87c:05015)

21. E. Sen, *The kind on Apostol-Euler polynomials of higher order arising from Euler basis*, Adv. Stud. Contemp. Math. (Kyungshang) **23** (2013), no. 2, 337–345.

DEPARTMENT OF MATHEMATICS, SOGANG UNIVERSITY, SEOUL 121-742, REPUBLIC OF KOREA
E-mail address: `dskim@sogang.ac.kr`

DEPARTMENT OF MATHEMATICS, KWANGWOON UNIVERSITY, SEOUL 139-701, REPUBLIC OF KOREA
E-mail address: `tkkim@kw.ac.kr`